

Broccoli

what's inside

- ▶ What's So Great about Broccoli?
- ▷ Selecting and Storing Broccoli
- **▷** Varieties of Broccoli
- **▷** Fitting Broccoli into MyPyramid
- ▶ Preparation Tips
- **▷** Recipe Collection
- □ Grow Your Own Broccoli





What's So Great about Broccoli?

Broccoli is an excellent source of vitamin C and dietary fiber. It's a good source of potassium and vitamin A. Broccoli is low in calories. It is also fat and cholesterol free. Broccoli contains the phytonutrient (plantnutrient) sulforaphane (pronounced: sul-for a-fain). Sulforaphane has been shown in some studies to reduce risk of cancer. Broccoli can be mixed with a variety of foods. It can be used as a soup, salad, flavoring for grains, or even as a main dish.

Selecting and Storing Broccoli

Broccoli is available all year. In some markets the florets (tops) of the broccoli are sold without the stalk.

Look for:

- Firm, tightly bunched florets
- Florets should be dark green, sage green, or even greenish purple
- Stems should not be too thick or too tough

Avoid

- Open buds or yellow flowers
- Yellowish green color
- Wilted or limp broccoli
- Soft, slippery, water-soaked spots on buds

Storage

Place unwashed broccoli in a plastic bag. Keep the broccoli in the vegetable crisper of the refrigerator. It will keep in the refrigerator for up to 4 days. Cooked leftovers may be refrigerated for 2 to 3 days in a tightly covered container.

Why is Vitamin C Important?

Broccoli is an excellent source of vitamin C. Vitamin C, also known as ascorbic acid, is needed for growth and repair of body tissue. Vitamin C helps the body make collagen,

a tissue needed for healthy bones, teeth, gums and blood vessels. Vitamin C is water soluble (dissolves in water) which means that you should have it every day, because your body does not store it.



Varieties of **Broccoli**



The most common type of broccoli sold in the U.S. is called sprouting, or Italian green, broccoli. Another name for this type of broccoli is Calabrese. This name came from the Italian province of Calabria where it was first grown. Sprouting broccoli has light-green stalks. Umbrella-shaped groups of purplishgreen florets top the stalks. Other broccoli varieties are:

<< Broccolini

This type is a cross between broccoli and Chinese kale.

Broccoflower >>

This type is a cross between broccoli and cauliflower. This broccoli comes in a single large head. Broccoflower is a light green, which is a mix of the white cauliflower and dark green broccoli.



<< Broccoli sprouts

Broccoli sprouts are a good source of sulforaphane. These tend to be expensive. Mix them with other vegetables to keep costs low.



Fitting Broccoli into MyPyramid

The **GREEN** triangle of MyPyramid is the vegetable section. Vegetables may be raw or cooked; fresh, frozen, or canned. Age, gender, and level of physical activity determine the daily veg-

etable need. For a 2000 calorie diet, it is recommended having about 2 ½ cups of vegetables daily. On MyPyramid, broccoli is a dark green vegetable. For a 2000 calorie diet, 3 cups of dark green vegetables are recomended per week. That's less than ½ cup per day! For more information on vegetable serving sizes visit www.mypyramid.gov.



Nutrition Facts 1 cup cooked Broccoli

Serving Size (148g) Servings Per Container Amount Per Serving Calories 40 Calories from Fat 5 Total Fat 1g Saturated Fat 0g 0% 0% Cholesterol 0mg Sodium 40mg 2% Total Carbohydrate 8g 3% Dietary Fiber 4g Sugars 3g Protein 4g Vitamin A 90% Vitamin C 230% Calcium 8% Iron 8% 'Percent Daily Values are based on a 2,000 calone diet. Your daily values may be higher or lower depending on your calorie needs

Preparation Tips

Boiling: Boiling broccoli creates a mild taste. Boiling can cause a loss of vitamin C and other nutrients. When boiling, bring the water to a rapid boil before adding the broccoli. Cooking time is 5 to 7 minutes.

Stir-frying: Stir-fry broccoli for two minutes. Add a little broth or water. Cover the pan. Let the broccoli steam until crisp-tender. Cooking time is 4 to 5 minutes.

Steaming: Steaming preserves the most nutrients. Steaming also keeps

the florets from breaking apart. Steam broccoli for 1 to 2 minutes. Then uncover the pot for 10 to 15 seconds. Uncovering the pot lets the gasses escape. Total cooking time is 5 to 7 minutes.

Microwaving: In a microwave-safe dish, arrange one pound of broccoli in a wheel-spoke fashion. The florets should point toward the center. Add 1/4 cup water and cover. Rotate the container halfway through the cooking time. Total cooking time is 6 to 10 minutes.



Recipe Collection

Creamy Broccoli Soup

Serves 4

INGREDIENTS

11/2 cups chicken broth

1/2 cup chopped onion

2 cups cut broccoli

1/2 teaspoon dried, crushed thyme

2 small bay leaves

2 Tablespoons margarine

2 Tablespoons flour

1/4 teaspoon salt

few dashes of pepper

1 cup nonfat milk

Garlic powder to taste

Estimated cost: Per Recipe: \$ 1.75 Per Serving: \$ 0.44

INSTRUCTIONS

1. In a saucepan combine chicken broth, chopped onion, broccoli, thyme, bay leaf and garlic powder.

Bring mixture to a boil. Reduce heat.
 Cover and simmer for 10 minutes or until vegetables are tender. Remove bay leaf.
 Place half of the mixture in a blender or food processor. Cover and blend 30 to 60 seconds or until smooth. Pour mixture into

a bowl. Repeat with remaining vegetable mixture. Set all aside.

4. In the same saucepan melt the margarine. Stir in flour, salt, and pepper. Add the milk all at once. Stir rapidly with a wire whisk.

5. Cook and stir until mixture is thickened and bubbly. Stir in the blended broccoli mixture. Cook and stir until soup is heated through. Season to taste with additional salt and pepper.

Options: Use butter in place of the margarine. Other types of milk (1%, 2%, whole) can be used in place of non-fat milk. These would change the nutrition information slightly. Chicken broth can be canned, boxed, or homemade.





Recipe Collection

Broccoli Salad

Makes 8 servings

INGREDIENTS

6 cups chopped broccoli

- 1 cup raisins
- 1 medium peeled and diced red onion
- 2 Tablespoons sugar
- 8 cooked and crumbled bacon
- slices (optional)
- 2 Tablespoons lemon juice
- 3/4 cup low fat mayonnaise

INSTRUCTIONS

- 1. Combine all ingredients in a medium bowl
- 2. Mix well
- 3. Chill for 1 to 2 hours
- 4. Serve

Options: Regular mayonnaise can be used in place of low fat mayonnaise. This would increase the fat grams. Bacon can be omitted from the recipe. The bacon adds to the total fat, cholesterol, protein, and sodium.

Estimated cost: Per Recipe: \$ 2.82 Per Serving: \$ 0.35



es per gram: Fat 9 · Carbohydrate 4 · Prosen 4

Broccoli Chicken Cheese Casserole

Makes 8 servings

INGREDIENTS

- 1 cup rice, uncooked
- 1 package (10 ounce) broccoli, frozen
- 3 cups chicken, cooked
- 2 Tablespoons margarine or butter
- 1/4 cup flour
- 2 cups chicken broth*
- 1/4 cup Parmesan cheese (optional)
- *Chicken broth can be canned, boxed, or homemade (from cooking the chicken).

Estimated cost: Per Recipe: \$8.97 Per Serving: \$1.12

INSTRUCTIONS

- 1. Cook rice in 2 cups of water or per the directions on the package.
- 2. Let broccoli thaw.
- 3. Make sure there are no bones with the
- chicken. Cut chicken into bite-sized pieces.
- 4. Melt butter in large sauce pan.
- 5. Add flour to melted margarine and stir. This will be lumpy.
- 6. Slowly add chicken broth to margarine/flour. Stir to remove lumps and thicken.
- 7. Add cheese and stir.
- 8. Add rice, broccoli, and chicken. Stir.9. Put in a casserole pan. Bake at 350°F for
- 25-30 minutes.







Grow Your Own **Broccoli**

Broccoli plants are often started indoors. Then broccoli is transplanted (moved) outdoors when the small plants are an inch or two tall.

Broccoli can survive through frost. But broccoli does not grow well in very hot weather. Plant broccoli in the spring or fall. For fall planting, start seedlings in midsummer. Transplant into the garden in late summer. Plant seeds 1/4 to 1/2 inch deep. Set transplants slightly deeper than they were grown. Plant 18 to 24 inches apart in the row. Space rows 36 inches apart. Broccoli plants grow tall, often reaching a height of 2 1/2 feet. Green buds develop first in one large central head. Later buds grow in several smaller side shoots.


After the central head of the broccoli has grown, cut it with 5 to 6 inches

of stem. Do this after the head is fully developed, but before it begins to loosen and separate. Removing the central head makes the sides hoots grow out from the lower leaves. Broccoli can be harvested for several weeks this way. For more information, contact your local Cooperative Extension Educator in the phone book or at www. extension.psu.edu/

ACTIVITY ALLEY

Broccoli Coloring Activity


Directions: Color the picture below.



ACTIVITY ALLEY

Broccoli Crossword Puzzle

Directions: Answer the clues to test how much you know about broccoli.



Across

- 3. In MyPyramid, broccoli is a _____vegetable. For a 2000 calorie diet, it is recommended that you eat about 3 cups of this colored vegetable per week.
- 5. Broccoli is an excellent source of this.
- 6. _____ variety of broccoli is a mix of broccoli and Chinese kale.
- 7. Avoid purchasing this color of broccoli.
- 8. Broccoli contains this phytonutrient. This phytonutrient has been found to help prevent cancer.
- 9. Broccoli _____ are a good source of sulforaphane.

Down

- 1. This variety of broccoli is most common in the US. It has light-green stalks. Umbrella-shaped groups of purplish-green florets top the stalks.
- 2. This variety is a cross between broccoli and cauliflower.
- 4. Another name for sprouting type of broccoli is ______.

References

Pennsylvania Nutrition Education Tracks, Pennsylvania State University, Cooperative Extension, State College, Pennsylvania, 2007.

P. Basiotis, A. Carlson, S. Gerrior, W. Juan, and M. Lino, The Healthy Eating Index 1999-2000, CNPP-12, Center for Nutrition Policy and Promotion, USDA, December 2002.

Prior, R. L. Fruits and Vegetables in the Prevention of Cellular Oxidative Damage, American Journal of Clinical Nutrition, 2003, 78, (Supplement) 570S-8S.

Nutrition for Everyone: Eat Your Fruits and Vegetables, Division of Nutrition, Physical Activity and Obesity, National Center for Chronic Disease Prevention and Health Promotion, Center for Disease Control and Prevention: May 2008.

2005 Dietary Guidelines for Americans, Center for Nutrition Policy and Promotion, US Department of Agriculture.

Krebs-Smith, S. M. and others. Fruits and Vegetable Intakes of Children and Adolescents in the United States, Archives of Pediatrics and Adolescent Medicine. Volume 50, Number 1, January 1996.

Krebs-Smith, S.W. and others. Choose a Variety of Fruits and Vegetables Daily: Understanding the Complexities, Journal of Nutrition. 2001; 131; 487S-501.

Gillman, H. Enjoy Your Fruits and Vegetables, British Medical Journal. 1966; 313, 765-766.


Hung, H.C. and others. Fruits and vegetables Intake and Risk of Major Chronic Disease, Journal of National Cancer Institute. 2004; 96: 1577-84.

Krinsky, N.I. Biologic Mechanisms of the Protective Roll of Luctein and Zeaxanthin in the Eyes, Annual Review of Nutrition. 2003; 23: 171-201.

Christen, W. G. and others. Fruit and Vegetable Intake and the Risk of Cataracts in Women, American Journal of Clinical Nutrition. 2005; 81: 1417-22



Cooperative Extension Service University of the District of Columbia 4200 Connecticut Ave. NW Washington, DC 20008 (202) 274-7115



This material was funded by USDA's Food Stamp Program. The Food Stamp Program provides nutrition assistance to people with low income. It can help buy nutritious foods for better diet. To find out more contact the Income Maintenance Administration @ 202-724-5506 to help identify how to get services. In cooperation with the U.S. Department of Agriculture and District of Columbia Government, Cooperative Extension Service and Agricultural Experiment Station programs and employment opportunities are available to all people regardless of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, marital status or family status. In accordance with Federal law and U.S. Department of Agriculture's policy, this institution is prohibited from discriminating on the basis of race, color, national origin, sex age, religion, political beliefs or disability. To file a complaint of discrimination, write USDA, Director of Civil Rights, Room 326-W, Whitten Building, 1400 Independence Avenue, SW, Washington, D.C. 20250 or call (202) 720-5694(voice and TDD). USDA is an equal opportunity provider and employer.